

Übungsblatt 3

Exercise 1. Let M be a smooth manifold and let $f: M \rightarrow \mathbb{R}^k$ be a smooth function. Show that $f \circ \varphi^{-1}: \varphi(U) \rightarrow \mathbb{R}^k$ is smooth for *any* chart (U, φ) for M .

Exercise 2. Let M, N, P be smooth manifolds. Show that the following hold.

- (1) Any constant map $c: M \rightarrow N$ is smooth.
- (2) The identity map $M \rightarrow M, p \mapsto p$, is smooth.
- (3) Let $U \subset M$ be an open subset and consider it as manifold (with the induced topology). Then the inclusion $U \hookrightarrow M$ is smooth.
- (4) If $F: M \rightarrow N$ and $G: N \rightarrow P$ are smooth, then $G \circ F: M \rightarrow P$ is smooth.

Exercise 3. Let \mathbb{B} be the unit sphere in \mathbb{R}^n . We consider the maps $F: \mathbb{B} \rightarrow \mathbb{R}^n$ and $G: \mathbb{R}^n \rightarrow \mathbb{B}$

$$F(x) = \frac{x}{\sqrt{1 - |x|^2}}, \quad G(y) = \frac{y}{\sqrt{1 + |y|^2}}$$

Show that F and G are smooth and inverse to each other.

Exercise 4. Let $f: M \rightarrow N$ be a smooth map between manifolds. Show that f is continuous and open.

Exercise 5. Let \mathbb{S}^1 be the unit circle in \mathbb{R}^2 . Show that for any integer $n \geq 1$, the map $\mathbb{S}^1 \rightarrow \mathbb{S}^1, z \mapsto z^n$, is smooth.

Exercise 6. Show that $\mathbb{R}\mathbb{P}^n$ is a smooth manifold and that the quotient map $\pi: \mathbb{R}^{n+1} \setminus \{0\} \rightarrow \mathbb{R}\mathbb{P}^n$ is smooth.